



SEEING

3 lectures on the physics of sight


- 1) *Light and imaging*
- 2) *The retina and colour vision*
- 3) *Measuring light and seeing more*

Dr John S. Reid
 Department of Physics
 University of Aberdeen



The Electromagnetic Spectrum

- Light is a tiny part of a huge spectrum of electromagnetic waves (section 24.2)



- Light in vacuum has **wavelengths** between ~400 nm and ~800 nm; 1 nm \equiv 10^{-9} m (a *nanometer*)
- 1 μ m \equiv 10^{-6} m (a *micron*)



A  of waves

Simple waves have 3 properties

- Wavelength** (λ in **m**) - how far before they repeat
- Frequency** (f in **Hz**) - the number of repetitions per second
- Speed** (v in **ms⁻¹**) - the speed of light in vacuum is always represented by the letter c
- The three quantities are simply related

$$\lambda = \frac{v}{f}$$



Example

- The human eye sees different frequencies as different colours. What frequency corresponds to **green light** of wavelength 550 nm in vacuum?
- Rearranging the relationship gives:

$$f = \frac{c}{\lambda}$$



- $c = 3.00 \times 10^8 \text{ ms}^{-1}$; 550 nm \equiv 5.50×10^{-7} m
- Therefore $f = 5.45 \times 10^{14}$ Hz




Electromagnetic?

- James Clerk Maxwell, former professor at Marischal College, Aberdeen, established the mathematical equations that describe electricity and magnetism
- He predicted that radiation would travel at speed c in vacuum and hence that light was an electromagnetic phenomenon



Hot body emission

- All bodies** emit electromagnetic radiation
- The spectral spread of this radiation is determined by the **temperature** of the body and a fundamental law of physics



Appearance of hot bodies

- Red is the colour of bodies that just glow; yellow, white and blue for really hot bodies
 - the concept of **colour temperature** is used by architects and others to label the spectrum of incident light they perceive

Emission from cooler bodies

- The spectrum of radiant energy was predicted by Planck
- Bodies at room temperature emit radiation in the far IR
- Man-made detectors can 'see' at other wavelengths than visible light

Seeing in the IR

- Meteorological satellites take both visible light and IR pictures
- Each picture element is called a *pixel*. A geostationary satellite takes pictures of at least 5000×5000 pixels in about 15 minutes
- The picture here was taken at midnight at a wavelength of about $6.5 \mu\text{m}$

How a picture is built up

- Meteosat forms its images by scanning a 'point detector', line by line over the scene below
- TV pictures are built up in the same way. The lines that build up the picture are called *raster lines*
- Our eye has an **area detector** at the rear, called the *retina*

A pin-hole eye

- Take an enclosure with a screen at the rear; make a tiny pin-hole and you have a *pin-hole camera*
- If the pin-hole is small ($\sim 0.3 \text{ mm}$) a very dim but respectable image appears on the screen
- The device relies on the property that **rays of light travel in straight lines**

An eye with a lens

- The lens collects a whole cone of light rays coming from each object point and focuses them down to one image point
- The function of the lens is to improve the illumination of the image
- If the lens fails to bring the image forming rays to a point, the image will be blurred


The iris



- The iris controls the aperture of the eye
- The smaller the aperture, the better quality the image
- The purpose of the iris is to give the eye the smallest aperture consistent with a bright enough image

13

Eye parts



- Cornea
- Lens
- Retina
- Optic nerve
- Humours
- Iris
- Pupil
- Fovea

14

Anatomy of the eye




- What the scalpel reveals

www.orlions.org/eyetour/fig24set.html

15

The bending of light

- To form an image, light rays must be bent
- The bending of light is known as **refraction**
- The law of refraction of light is *Snell's law*
- The ability of a medium to refract light is given by its *refractive index, n*



$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$

16

Numerical example of bending

- Light strikes the cornea at an angle of incidence (θ_1) of 20° . The cornea has a refractive index of 1.376. What angle (θ_2) does the light leave the cornea surface?
- Take $n_1 = 1.000$

$$1.000 \sin(20^\circ) = 1.376 \sin \theta_2$$

$$\therefore \sin \theta_2 = (0.342)/1.376 = 0.249$$


$$\therefore \theta_2 = 14.4^\circ$$

- The light leaves at 14.4° to the surface normal

17

Imaging by the eye


- 75% of the imaging power of your eye is at the front of the cornea. Look after it!
- The lens is an adjustable element for focusing objects at different distances
- The variable focusing of the eye is known as its *accommodation*



18



Imaging by a convex lens



- Broadly speaking, the eye images like a powerful convex lens
- The images of all objects we can see are small, inverted and form on the retina
- In the **next lecture**, we shall look at properties of the retina

19

